



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 211/74, A61K 31/395, C07D 205/085, 223/12, 207/273, A61K 31/445, 31/40		A1	(11) International Publication Number: WO 97/05111 (43) International Publication Date: 13 February 1997 (13.02.97)
(21) International Application Number: PCT/EP96/02962 (22) International Filing Date: 5 July 1996 (05.07.96) (30) Priority Data: 9515411.8 27 July 1995 (27.07.95) GB (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN S.P.A. [IT/IT]; Via Robert Koch, 1,2, I-20152 Milan (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): PEVARELLO, Paolo [IT/IT]; Piazza San Pietro in Ciel d'Oro, 7/A, I-27100 Pavia (IT). AMICI, Raffaella [IT/IT]; Via N. Rocca, 11, I-29100 Piacenza (IT). VARASI, Mario [IT/IT]; Via Giambellino, 80, I-20146 Milan (IT). MAI, Roberto [IT/IT]; Via Bossi, 24, I-21047 Saronno (IT). SALVATI, Patricia [IT/IT]; Via Valese, 16/C, I-20020 Arese (IT). (74) Common Representative: PHARMACIA & UPJOHN S.P.A.: Patent and Trademark Dept., Viale Pasteur, 10, I-20014 Nerviano (IT).		(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>	
(54) Title: N-(4-SUBSTITUTED-BENZYL)-2-AMINOLACTAM DERIVATIVES			
<p style="text-align: right;">(I)</p>			
(57) Abstract			
<p>Novel N-(4-substituted-benzyl)-2-aminolactams compounds, having antiepileptic, anti-Parkinson, neuroprotective, anti-depressant, antispasmodic and/or hypnotic activity, of general formula (I) wherein: m is zero, 1, 2 or 3; n is zero, 1, 2 or 3; X is -O-, -S-, -CH2- or -NH-; each of R and R₁ independently is hydrogen, C₁-C₈ alkyl, halogen, hydroxy, C₁-C₄ alkoxy or trifluoromethyl; each of R₂, R₃ and R₄ independently is hydrogen, C₁-C₈ alkyl optionally substituted by a hydroxy group, or C₃-C₇ cycloalkyl; and the pharmaceutically acceptable salts thereof, are disclosed.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

N-(4-SUBSTITUTED-BENZYL)-2-AMINOLACTAM DERIVATIVES

The present invention relates to novel N-(4-substituted-benzyl)-2-aminolactams, to their use as therapeutic agents, to a process for their preparation and to pharmaceutically compositions containing them.

It has been found that novel N-(4-substituted-benzyl)-2-aminolactam derivatives as herein defined have valuable biological properties, in particular as antiepileptic, anti-Parkinson, neuroprotective, anti-depressant, antispastic and/or hypnotic agent.

The present invention provides novel compounds of formula (I)



wherein:

m is zero, 1, 2 or 3;

n is zero, 1, 2 or 3;

X is -O-, -S-, -CH₂- or -NH-;

each of R and R₁ independently is hydrogen, C₁-C₆ alkyl, halogen, hydroxy, C₁-C₄ alkoxy or trifluoromethyl;

each of R₂, R₃ and R₄ independently is hydrogen, C₁-C₆ alkyl optionally substituted by a hydroxy group, or C₃-C₇ cycloalkyl;

and the pharmaceutically acceptable salts thereof.

The pharmaceutically acceptable salts of the compounds of formula (I) include acid addition salts with inorganic, e.g. hydrochloric, hydrobromic, sulphuric, and phosphoric acids,

or organic, e.g. acetic, propionic, lactic, oxalic, malic, maleic, tartaric, citric, benzoic, mandelic, salicylic, C₁-C₄ alkylsulfonic and fumaric acids.

- 5 The compounds of the formula (I), their pharmaceutically acceptable salts may also form pharmaceutically acceptable solvates, such as hydrates, which are also object of the present invention.

The alkyl and alkoxy groups may be branched or straight groups.

- 10 A C₁-C₆ alkyl group is preferably a C₁-C₄ alkyl group, in particular methyl, ethyl, *n*- and *iso*-propyl, *n*-, *iso*-, *sec*- and *tert*-butyl, more preferably methyl or ethyl.

A C₁-C₆ alkyl group substituted by hydroxy is typically hydroxymethyl.

- 15 Representative examples of C₁-C₄ alkoxy groups include methoxy or ethoxy.

A halogen atom is e.g. chlorine, fluorine, bromine, in particular chlorine and fluorine, more preferably fluorine.

- A C₃-C₆ cycloalkyl group is, for instance, a cyclopropyl, 20 cyclohexyl or cycloheptyl group, in particular cyclopropyl.

The present invention also include within its scope both the metabolites and the pharmaceutically acceptable bioprecursors (otherwise known as pro-drugs) of the compounds of formula (I).

- 25 Compounds of formula (I) contain an asymmetric carbon atom and have optical *l* and *d* isomers. Accordingly, the invention includes all the possible isomers, in particular *l* and *d* isomers and their mixtures.

- 30 Preferred compounds of the invention are the compounds of formula (I) wherein

m is 1 or 2;

- n is 1 or 2;
X is -O-, -S- or -NH-;
R is hydrogen;
R₁ is hydrogen or halogen;
5 each of R₂ and R₄ independently is hydrogen or C₁-C₄ alkyl;
R₃ is hydrogen or C₁-C₄ alkyl optionally substituted by a hydroxy group; and the pharmaceutically acceptable salts thereof.
- 10 More preferred compounds of the invention are the compounds of formula (I), wherein
m is 1;
n is 1;
X is -O- or -NH-;
15 R₁ is hydrogen or halogen;
R₂ is hydrogen or C₁-C₄ alkyl;
R₃ is hydrogen or C₁-C₄ alkyl optionally substituted by hydroxy;
R and R₄ are hydrogen; and the pharmaceutically acceptable
20 salts thereof.

Examples of specific compounds of the invention are:


- 3-[4-(3-fluorobenzyloxy)benzylamino]-pyrrolidin-2-one;
3-[4-(3-chlorobenzyloxy)benzylamino]-pyrrolidin-2-one;
25 3-[4-(4-chlorobenzyloxy)benzylamino]-pyrrolidin-2-one;
3-[4-(3-bromobenzyloxy)benzylamino]-pyrrolidin-2-one;
3-[4-(4-fluorobenzyloxy)benzylamino]-pyrrolidin-2-one;
3-[4-(2-fluorobenzyloxy)benzylamino]-pyrrolidin-2-one;
3-[4-(3-fluorobenzyloxy)benzylamino]-azetidin-2-one;
30 3-[4-(3-fluorobenzyloxy)benzylamino]-piperidin-2-one;
3-[4-(3-fluorobenzyloxy)benzylamino]-azepan-2-one;

- 3-[4-(3-fluorobenzylamino)benzylamino]-pyrrolidin-2-one;
 3-[4-(benzylsulfanyl)benzylamino]-pyrrolidin-2-one;
 3-[[4-(3-fluorobenzyloxy)benzyl]methylamino]-pyrrolidin-2-one;
 5 3-[[4-(3-fluorobenzyloxy)benzyl]methylamino]-1-hydroxymethyl-pyrrolidin-2-one;
 3-[4-(3-fluorobenzyloxy)benzylamino]-1-methyl-pyrrolidin-2-one;
 3-[[4-(3-chlorobenzyloxy)benzyl]methylamino]-pyrrolidin-2-one;
 10 one;
 3-[[4-(3-bromobenzyloxy)benzyl]methylamino]-pyrrolidin-2-one,
 if the case, either as a single isomer or as a mixture of isomers thereof, and the pharmaceutically acceptable salts thereof.

15


The compounds of the invention and the salts thereof can be obtained by a process comprising:

- a) reaction of a compound of formula (II)



20

wherein n, R, R₁ and X are as defined above, with a compound of formula (III)




25


wherein m, R₃ and R₄ are as defined above, thus obtaining a compound of formula (I) in which R₂ is

hydrogen; or

b) reacting a compound of formula (IV)



5 wherein R , R_1 , R_3 , R_4 , m , n and X are as defined above,
with a compound of formula (V) or (VI)



wherein W is a halogen atom; R'_2 is a C_1 - C_4 alkyl and
 R''_2 is hydrogen or C_1 - C_3 alkyl, thus obtaining a
 10 compound of the invention in which R_2 is C_1 - C_4 alkyl;
 and, if desired, converting a compound of the invention
 into another compound of the invention and/or, if
 desired, converting a compound of the invention into a
 pharmaceutically acceptable salt and/or, if desired,
 15 converting a salt into a free compound and/or, if
 desired, separating a mixture of isomers of compounds of
 the invention into a single isomer.

20 All the processes described hereabove are analogy processes
 and can be carried out according to well known methods in
 organic chemistry.

A compound of formula (IV) is a compound of formula (I) in
 which R_2 is hydrogen.

25 The reaction of a compound of formula (II) with a compound of
 formula (III) to give a compound of formula (I) or (IV) is a
 reductive amination reaction which can be carried out
 according to well known methods. According to a preferred
 embodiment of the invention it may be performed under

nitrogen atmosphere, in a suitable organic solvent, such as an alcohol, e.g. a lower alkanol, in particular methanol, or in acetonitrile, at a temperature ranging from about 0°C to about 40°C, in the presence of a reducing agent, the most appropriate being sodium cyanoborohydride. Occasionally molecular sieves can be added to the reaction mixture for facilitating the reaction.

In a compound of formula (V) the halogen W is preferably iodine. The alkylation reaction of a compound of formula (IV) with a compound of formula (V) can be carried out in a suitable organic solvent, such as an alcohol, e.g. methanol, ethanol or isopropanol, in particular in ethanol, at a temperature ranging from about 0°C to about 50°C.

The alkylation reaction of a compound of formula (IV) with an aldehyde of formula (VI) can be carried out in a suitable organic solvent, such as an alcohol, e.g. methanol, ethanol or acetonitrile in the presence of a suitable reducing agent, such as sodium cyanoborohydride, at a temperature ranging from about 0°C to about 30°C.

A compound of the invention can be converted, as stated above, into another compound of the invention by known methods. Process-variant b) above may be regarded as an example of optional conversion of a compound of the invention into another compound of the invention.

An isomer, e.g., a d- or l-isomer of a compound of the invention can be separately synthesized from optically pure starting material or separated from a racemate in a conventional manner.

Also the optional salification of a compound of the invention as well as the conversion of a salt into the free compound may be carried out by conventional methods.

When in the compounds of the present invention and in the

intermediate-products thereof, groups are present, which need to be protected before submitting them to the hereabove illustrated reactions, they may be protected before being reacted and then deprotected according to methods well known in organic chemistry.

The compounds of formula (II), (III), (V) and (VI) are known compounds or can be obtained by known methods.

PHARMACOLOGY

The compounds of the invention are active on the central nervous system (CNS) and can be used in therapy, for example as antiepileptics, in the treatment of Parkinson's disease and as neuroprotective agents, e.g. preventing or treating neuronal loss associated with stroke, ischemia, CNS trauma, hypoglycaemia or surgery and in treating or preventing neurodegenerative diseases such as Alzheimer's disease, amyotrophic lateral sclerosis, Down's syndrome or Huntington's disease; they can also be used as antidepressants, hypnotics and antispastic agents. The activity on the CNS of the compounds of the invention was evaluated on the basis of pharmacological methods, such as, for example, the antagonism of convulsions and lethality induced by intravenous injection of bicuculline in mice (Antiepileptic Drugs, D.M. Woodbury et al. eds., 2nd edition, Raven Press, New York, 1982), or the antagonism of maximal electroshock seizures (MES) (Woodbury, L.A. and Davenport, V.D., Arch. Int. Pharmacodyn. Ther. 92; 97-104, 1952).

A representative group of compounds of the invention was tested in the MES test as described before and found to be active as reported in Table 1 below.

Table 1

Compound	MES ED ₅₀
FCE 28861	23.7
FCE 29530	27.1
FCE 29487A	30.1
FCE 28860A	20.8
FCE 29481A	10.4
FCE 29483A	9.95
FCE 29495A	27.6
FCE 29532A	43.0
FCE 29531	27.3
FCE 29642A	9.82
FCE 29643	40.2
FCE 29646A	24.2
FCE 29823A	15.6
FCE 29821A	13.77

where:

- FCE 28861 means (S)-3-(4-(3-fluorobenzyloxy)benzylamino) piperidin-2-one;
- 5 FCE 29530 means (S)-3-(4-(3-fluorobenzyloxy)benzylamino) azetidin-2-one;
- FCE 29487A means (S)-3-(4-(3-fluorobenzyloxy)benzylamino) azepan-2-one; methanesulfonate;
- FCE 28860A (S)-3-(4-(3-fluorobenzyloxy)benzylamino) 10 pyrrolidin-2-one, methanesulfonate;
- FCE 29481A means (S)-3-(4-(3-bromobenzyloxy)benzylamino) pyrrolidin-2-one, methanesulfonate;
- FCE 29483A means (S)-3-(4-(2-fluorobenzyloxy)benzylamino) pyrrolidin-2-one, methanesulfonate;
- 15 FCE 29532A means (S)-3-(4-(benzylsulfanyl)benzylamino) pyrrolidin-2-one, methanesulfonate;

FCE 29531 means (S)-3-((4-(3-fluorobenzyloxy)benzyl)methylamino)-1-hydroxymethyl-pyrrolidin-2-one;

FCE 29642A means (S)-3-(4-(3-chlorobenzyloxy)benzylamino)pyrrolidin-2-one, methanesulfonate;

5 FCE 29643A (S)-3-((4-(3-fluorobenzyloxy)benzyl)methylamino)pyrrolidin-2-one, methanesulfonate;

FCE 29646A means (S)-3-(4-(4-fluorobenzyloxy)benzylamino)pyrrolidin-2-one, methanesulfonate;

FCE 29823A means (S)-3-(4-(3-fluorobenzyloxy)benzylamino)-1-
10 methyl-pyrrolidin-2-one, hydrochloride;

FCE 29821A means (S)-3-(4-(3-fluorobenzylamino)benzylamino)pyrrolidin-2-one, methanesulfonate;

ED₅₀ means effective dose (mg/Kg) in 50% of treated animals (mice) 1hour after per os (po) administration.

15

A patient is treated according to the present invention by a method comprising administering to the patient an effective amount of one of the compounds of the invention. In this way the present compounds can be used to treat disorders of the
20 central nervous system, for example epilepsy or Parkinson's disease; or as neuroprotective agents, anti-depressants, hypnotics or anti-spastic agents. The condition of a patient may thus be improved.

The compounds of the invention can be administered in a
25 variety of dosage forms, e.g. orally, in the form of tablets, capsules, sugar or film coated tablets, liquid solutions or suspensions; rectally in the form of suppositories; parenterally, e.g. intramuscularly, or by intravenous injection or infusion.

30 The dosage depends on the age, weight, conditions of the patient and on the administration route; for example, the dosage adopted for oral administration to adult humans e.g.

for the representative compound of the invention 3-[4-(3-chlorobenzoyloxy)benzylamino]-pyrrolidin-2-one may range from about 1 to about 500 mg pro dose, from 1 to 5 times daily.

The invention includes pharmaceutical compositions comprising
5 a compound of the invention in association with a pharmaceutically acceptable excipient (which can be a carrier or a diluent).

The pharmaceutical compositions containing the compounds of the invention are usually prepared following conventional
10 methods and are administered in a pharmaceutically suitable form.

For example, the solid oral forms may contain, together with the active compound, diluents, e.g. lactose, destrose, saccharose, cellulose, corn starch or potato starch;
15 lubricants, e.g. silica, talc, stearic acid, magnesium or calcium stearate, and/or polyethylene glycols; binding agents, e.g. starches, arabic gums, gelatin, methylcellulose, carboxymethylcellulose or polyvinyl pyrrolidone; disaggregating agents, e.g. a starch, alginic acid, alginates
20 or sodium starch glycolate; effervescing mixtures; dyestuffs; sweeteners; wetting agents such as lecithin, polysorbates, laurylsulphates; and, in general, non-toxic and pharmacologically inactive substances used in pharmaceutical formulations. Said pharmaceutical preparations may be
25 manufactured in known manner, for example, by means of mixing, granulating, tableting, sugar-coating, or film-coating processes.

The liquid dispersion for oral administration may be e.g. syrups, emulsions and suspension.

30 The syrups may contain as carrier, for example, saccharose or saccharose with glycerine and/or mannitol and/or sorbitol.

The suspension and the emulsion may contain as carrier, for

example, a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl alcohol.

The suspension or solutions for intramuscular injections may contain, together with the active compound, a pharmaceutically acceptable carrier, e.g. sterile water, olive oil, ethyl oleate, glycols, e.g. propylene glycol, and, if desired, a suitable amount of lidocaine hydrochloride. The solutions for intravenous injections or infusion may contain as carrier, for example, sterile water or preferably they may be in the form of sterile, aqueous, isotonic saline solutions.

The suppositories may contain together with the active compound a pharmaceutically acceptable carrier, e.g. cocoa butter, polyethylene glycol, a polyoxyethylene sorbitan fatty acid ester surfactant or lecithin.

The following examples illustrate but do not limit the invention.

Example 1

(S)-3-[4-(3-fluorobenzoyloxy)benzylamino]-pyrrolidin-2-one, methanesulfonate

To a solution of (S)-3-amino-2-pyrrolidinone hydrochloride (Synthesis, 1978, 614) (2.0 g; 0.0146 mol) in anhydrous methanol (60 ml), under stirring and nitrogen, 2.0 g of 3Å molecular sieves were added and then, in a single portion, NaBH_3CN (0.77 g; 0.0122 mol); after 10 minutes, 3.4 g (0.0148 mol) of 4-(3-fluorobenzoyloxy)benzaldehyde were added, in 40 ml of anhydrous methanol. After three hours the reaction was completed, the mixture filtered, the solution was evaporated to give a residue which was directly flash-chromatographed on

silica gel (eluent: CHCl_3 , 97: CH_3OH 3: 30% NH_4OH 0.3) to give a white solid (3.4 g; 74%). The free base thus obtained was treated with a stoichiometric amount of methanesulfonic acid to yield the title compound (m.p. 140.5-143.5°C; $[\alpha]_D^{25} +35.2$
5 (c=1.22, AcOH).

Analogously the following products can be obtained, starting from the corresponding aldehyde and the appropriate 3-aminolactam:

10 (S)-3-[4-(3-chlorobenzoyloxy)benzylamino]-pyrrolidin-2-one, methanesulfonate:

m.p. 138-144°C; $[\alpha]_D^{25} = -32.6$ (c=1.2, AcOH);

(S)-3-[4-(3-fluorobenzoyloxy)benzylamino]-piperidin-2-one:

15 m.p. 98-101°C; $[\alpha]_D^{25} = -3.7$ (c=1.0, AcOH);

(S)-3-[4-(3-bromobenzoyloxy)benzylamino]-pyrrolidin-2-one,

methanesulfonate: m.p. 138.5-143.5°C; $[\alpha]_D^{25} = -30.0$ (c=1, AcOH);

20 (S)-3-[4-(4-fluorobenzoyloxy)benzylamino]-pyrrolidin-2-one, methanesulfonate:

m.p. 141°C; $[\alpha]_D^{25} = -34.4$ (c=1.2, AcOH);

(S)-3-[4-(4-chlorobenzoyloxy)benzylamino]-pyrrolidin-2-one:

25 m.p. 172-176°C; $[\alpha]_D^{25} = -35.0$ (c=1, AcOH);

(S)-3-[4-(2-fluorobenzoyloxy)benzylamino]-pyrrolidin-2-one,

methanesulfonate: m.p. 176.5-179°C; $[\alpha]_D^{25} = -34.0$ (c=1, ACOH);

3-[4-(3-fluorobenzyloxy)benzylamino]-azetidin-2-one:

m.p. 89-91°C (dec.);

5

(S)-3-[4-(3-fluorobenzyloxy)benzylamino]-azepan-2-one,

methanesulfonate: m.p. 214-217.5°C; $[\alpha]_D^{25} = -13.1$ (c=1, H₂O);

(S)-3-[4-(3-fluorobenzylamino)benzylamino]-pyrrolidin-2-one,

10 dihydrochloride: m.p. 165°C; $[\alpha]_D^{25} = -30.3$ (c=1, ACOH);

(S)-3-[4-(benzylsulfanyl)benzylamino]-pyrrolidin-2-one,

methanesulfonate:

m.p. 138°C (dec.); $[\alpha]_D^{25} = -37.2$ (c=1.1, ACOH);

15

3-[4-(3-fluorobenzyloxy)benzylamino]-3-methyl-pyrrolidin-2-one, methanesulfonate; and

(S)-3-[4-(3-fluorobenzyloxy)benzylamino]-1-methyl-pyrrolidin-

20 2-one, hydrochloride:

m.p. 168-170°C; $[\alpha]_D^{25} = -31.6$ (c=1.2, NCOH).

Example 2

(S)-3-[[4-(3-fluorobenzyloxy)benzyl]methylamino]-pyrrolidin-2-

25 one

One g (0.00318 mol) of (S)-3-[4-(3-fluorobenzyloxy)benzylamino]pyrrolidin-2-one was dissolved in acetonitrile

(50 ml) under a nitrogen stream. To this mixture, 1.6 ml (0.0196 mol) of 37% formaldehyde and 0.29 g (0.00460 mol) of sodium cyanoborohydride were added at room temperature. After 20 minutes, glacial acetic acid was dropped up to neutrality of the solution. The mixture was stirred for 40 minutes and was evaporated to dryness. 40 ml of 2N KOH were added to the residue. After extracting with ethyl acetate, washing with N/2 KOH and then with water and brine, the organic layer was dried on Na_2SO_4 , then filtered and evaporated to obtain a residue which was flash-chromatographed on silica gel (eluant: CHCl_3 200: CH_3OH 3: 30% NH_4OH 0.2) to give 0.73 g (70%) of a white solid.

m.p. 89-92°C; $[\alpha]_D^{25} = -39.8$ (c=1.1, ACOH).

Analogously the following products can be prepared starting from the corresponding secondary amine:

3-{{[4-(3-chlorobenzoyloxy)benzyl]methylamino}-pyrrolidin-2-one; and

3-{{[4-(3-bromobenzoyloxy)benzyl]methylamino}-pyrrolidin-2-one.

20

Example 3

(S)-(+)-3-{{[4-(3-fluorobenzoyloxy)benzyl]methylamino}-1-hydroxymethyl-pyrrolidin-2-one

To a stirred solution of 1.45 g (0.00461 mol) of (S)-3-[4-(3-fluorobenzoyloxy)benzylamino]pyrrolidin-2-one and 3.7 ml (0.0493 mol) of 40% aqueous formaldehyde in 70 ml of acetonitrile was added 0.36 g (0.00573 mol) of sodium cyanoborohydride. The reaction mixture was stirred for 20 min, and then glacial acetic acid was added dropwise up to neutrality of the solution. Stirring was continued for an

additional 40 min, the solvent was evaporated at reduced pressure, and 40 ml of 0.5 N NaOH was added to the residue. After extracting with ethyl acetate and washing with brine, the organic layer was dried on Na_2SO_4 , then filtered and
5 evaporated to obtain a residue which was flash-chromatographed on silica gel (eluant: CHCl_3 90: MeOH 10) to give an oil. Treatment of the oil thus obtained with diisopropyl ether afforded the title compound as a white solid (0.95 g; 57%).

10 m.p. 96-98°C; $[\alpha]_D^{25} = +2.4$ (c=1.18, MeOH).


Example 4

With the usual methods of pharmaceutical technique, preparation can be made of capsules having the following
15 composition:

(S)-3-[4-(3-chlorobenzoyloxy)benzylamino]-pyrrolidin-2-one, methanesulfonate	50 mg
Talc powder	2 mg
Corn starch	2 mg
20 Microcrystalline cellulose	6 mg
Magnesium stearate	1 mg

CLAIMS

1. A compound of formula (I)



5 wherein:

m is zero, 1, 2 or 3;

n is zero, 1, 2 or 3;

X is -O-, -S-, -CH₂- or -NH-;

- each of R and R₁ independently is hydrogen, C₁-C₆ alkyl,
 10 halogen, hydroxy, C₁-C₄ alkoxy or trifluoromethyl;
 each of R₂, R₃ and R₄ independently is hydrogen, C₁-C₆ alkyl
 optionally substituted by a hydroxy group, or C₃-C₇
 cycloalkyl; or a pharmaceutically acceptable salt thereof.

15 2. A compound of formula (I), according to claim 1,
 wherein:

m is 1 or 2;

n is 1 or 2;

X is -O-, -S- or -NH-;

20 R is hydrogen;

R₁ is hydrogen or halogen;

each of R₂ and R₄ independently is hydrogen or C₁-C₄ alkyl;

R₃ is hydrogen or C₁-C₄ alkyl optionally substituted by a
 hydroxy group; or a pharmaceutically acceptable salt thereof.

25

3. A compound of formula (I), according to claim 1,
 wherein:

m is 1;

n is 1;

X is -O- or -NH-;

R₁ is hydrogen or halogen

5 R₂ is hydrogen or C₁-C₄ alkyl;

R₃ is hydrogen or C₁-C₄ alkyl optionally substituted by hydroxy;

R and R₄ are hydrogen; or a pharmaceutically acceptable salt thereof.

10

4. A compound selected from the group consisting of:

3-[4-(3-fluorobenzyloxy)benzylamino]-pyrrolidin-2-one;

3-[4-(3-chlorobenzyloxy)benzylamino]-pyrrolidin-2-one;

3-[4-(4-chlorobenzyloxy)benzylamino]-pyrrolidin-2-one;

15 3-[4-(3-bromobenzyloxy)benzylamino]-pyrrolidin-2-one;

3-[4-(4-fluorobenzyloxy)benzylamino]-pyrrolidin-2-one;

3-[4-(2-fluorobenzyloxy)benzylamino]-pyrrolidin-2-one;

3-[4-(3-fluorobenzyloxy)benzylamino]-azetidin-2-one;

3-[4-(3-fluorobenzyloxy)benzylamino]-piperidin-2-one;

20 3-[4-(3-fluorobenzyloxy)benzylamino]-azepan-2-one;

3-[4-(3-fluorobenzylamino)benzylamino]-pyrrolidin-2-one;

3-[4-(benzylsulfanyl)benzylamino]-pyrrolidin-2-one;

3-[[4-(3-fluorobenzyloxy)benzyl]methylamino]-pyrrolidin-2-one;

25 3-[[4-(3-fluorobenzyloxy)benzyl]methylamino]-1-hydroxymethylpyrrolidin-2-one;

3-[4-(3-fluorobenzyloxy)benzylamino]-1-methylpyrrolidin-2-one;

3-[[4-(3-chlorobenzyloxy)benzyl]methylamino]-pyrrolidin-2-


30 one;

3-([4-(3-bromobenzyloxy)benzyl]methylamino)-pyrrolidin-2-one, if the case, either as a single isomer or as a mixture of isomers thereof, and the pharmaceutically acceptable salts thereof.

5


5. A process for the preparation of a compound of formula (I), as defined in claim 1, or a pharmaceutically acceptable salt thereof, the process comprising:

a) reaction of a compound of formula (II)




10

wherein n, R, R₁ and X are as defined in claim 1, with a compound of formula (III)




wherein m, R₃ and R₄ are as defined in claim 1, thus
15 obtaining a compound of formula (I) in which R₂ is hydrogen;
or

b) reacting a compound of formula (IV)



wherein R, R₁, R₃, R₄, m, n and X are as defined in claim 1,
20 with a compound of formula (V) or (VI)



wherein W is a halogen atom; R'₂ is a C₁-C₄ alkyl and R''₂ is hydrogen or C₁-C₃ alkyl, thus obtaining a compound of the invention in which R₂ is C₁-C₄ alkyl; and, if desired, converting a compound of the invention into another compound of the invention and/or, if desired, converting a compound of the invention into a pharmaceutically acceptable salt and/or, if desired, converting a salt into a free compound and/or, if desired, separating a mixture of isomers of compounds of the invention into a single isomer.

10

6. A pharmaceutical composition containing a suitable carrier and/or diluent and, as an active principle, a compound of formula (I), as defined in claim 1, or a pharmaceutically acceptable salt thereof.

15

7. A compound of formula (I), as defined in claim 1, or a pharmaceutically acceptable salt thereof, for use as an active therapeutic substance.

20

8. A compound of formula (I), as defined in claim 1, or a pharmaceutically acceptable salt thereof, for use as antiepileptic, anti-Parkinson, neuroprotective, antidepressant, antispastic and hypnotic agent, and in treating and preventing neurodegenerative diseases.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 96/02962

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D211/74 A61K31/395 C07D205/085 C07D223/12 C07D207/273
A61K31/445 A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP, A, 0 362 941 (MERCK SHARP & DOHME LTD.) 11 April 1990 see the whole document ---	1-8
A	EP, A, 0 171 159 (YAMANOUCHI PHARMACEUTICAL CO. LTD.) 12 February 1986 see the whole document ---	1-8
A	EP, A, 0 001 601 (TAKEDA YAKUHIIN KOGYO KABUSHIKI KAISHA) 2 May 1979 see the whole document, especially pages 1-2 -----	1-8



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

'A': document member of the same patent family

Date of the actual completion of the international search

17 October 1996

Date of mailing of the international search report

8.11.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Scruton-Evans, I

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No

PCT/EP 96/02962

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-362941	11-04-90	AT-T- 131811	15-01-96
		CA-A- 1337770	19-12-95
		DE-D- 68925173	01-02-96
		DE-T- 68925173	29-08-96
		ES-T- 2081834	16-03-96
		JP-A- 2275851	09-11-90
		US-A- 4925867	15-05-90

EP-A-171159	12-02-86	JP-A- 61010578	18-01-86
		JP-A- 61027960	07-02-86
		JP-B- 5033716	20-05-93
		JP-A- 61078785	22-04-86
		JP-B- 5033717	20-05-93
		JP-A- 61078786	22-04-86
		CA-A- 1270097	05-06-90
US-A- 4719207	12-01-88		

EP-A-1601	02-05-79	JP-C- 1436840	25-04-88
		JP-A- 55015432	02-02-80
		JP-B- 62045863	29-09-87
		JP-A- 54055562	02-05-79
		AU-B- 518986	29-10-81
		AU-A- 4024278	03-04-80
		CA-A- 1102338	02-06-81
US-A- 4198514	15-04-80		
